



CIRCAID
by medi

VALCO
800.361.3153
www.valco.ca

RX PHYSICIAN'S
PRESCRIPTION

Graduated Medical Compression Stockings

Date: _____

Patient Name: _____

Diagnosis: _____

Physician Signature: _____ **(Dispense as written)
no substitution**

Practice #: _____ Dispense Quantity: _____ Rep: _____

Please indicate Compression and Style for Compression garment.

RX COMPRESSION (select one)

- | | | |
|--|--|---|
| <input type="checkbox"/> Preventive | <input type="checkbox"/> Class I: 20-30 mmHg | <input type="checkbox"/> Class II: 30-40 mmHg |
| <input type="checkbox"/> Anti-embolism
(Non ambulatory) | <input type="checkbox"/> Class III: 40-50 mmHg | <input type="checkbox"/> Class IV: 50-60 mmHg |

NON ELASTIC COMPRESSION

- CircAid Juxtalite® Venous Maternity Belt
 CircAid Juxtafit® Lymphedema

PNEUMATIC COMPRESSION

- 3 12
 Venous Lymphedema

(select one style form chart)

STYLE



Calf

Thigh

Panty

**Lymphedema
sleeve**

Glove

Left

Right

Left

Right

Dealer: _____

PHYSICIAN'S THERAPY GUIDE

INDICATIONS

SUPPORT

- Slight varicosities
- Tired aching legs

RX COMPRESSION

CLASS I: 20-30 mmHg

- Heaviness & fatigue in legs
- Minor varicosities
- Initial varices during pregnancy
- Post surgical
- Minor ankle, leg & foot swelling

- Post sclerotherapy
- Superficial thrombophlebitis
- Prophylactic treatment during pregnancy
- Hereditary disposition toward vein disease

Recommendations

mediven elegance®
mediven® for men
mediven® active
mediven® comfort
mediven plus®
juxtalite®

CLASS II: 30-40 mmHg

- Moderate varicosities
- Moderate edema
- Lymphatic edema
- Management of venous ulcers
- Prevention of ulcer recurrence
- Prevention of post-thrombotic syndrome

- Orthostatic hypotension
- Post surgical
- Post sclerotherapy
- Effective scar formation after burns
- Pregnancy with previous phlebitis
- Stasis dermatitis due to CVI

Recommendations

mediven elegance®
mediven® for men
mediven® active
mediven® comfort
mediven plus®
mediven® forte
mediven ulcer kit®
juxtalite®
juxtafit®

CLASS III: 40-50 mmHg

- Severe varicosities
- Severe edema
- Lymphatic edema
- Management of venous ulcers
- Post phlebotic syndrome

- Chronic venous insufficiency
- Resersible lymphedema
- Venous ulcers
- Post surgery
- Orthostatic hypotension

Recommendations

mediven plus®
mediven® forte
juxtafit®

CONTRAINDICATIONS:

According to the guideline of the German Phlebological Society:

Absolute Contraindications: Advanced peripheral arterial occlusive disease, decompensated heart disease, septic phlebitis, phlegmasia coerulea dolens, congestive heart failure, cutaneous infections, severe arterial insufficiency. **Relative contraindications:** suppurative dermatoses, intolerance of compression stocking fabric, sensory disturbances of the limb, advanced peripheral neuropathy, primary chronic arthritis.

J. Vaillancourt Corp./Medi assumes no liability as a result of any contraindicated use of this product.